

Near Horizon Geometry of Extreme Black Holes and Colliding Waves

M. Halilsoy

Physics Department., Eastern Mediterranean University G.Magosa, North

Cyprus, Mersin 10, Turkey

email: mustafa.halilsoy@emu.edu.tr

(February 7, 2008)

arXiv:gr-qc/0210055v1 17 Oct 2002

Typeset using REVTeX

Abstract

The correspondence between black holes and colliding waves extends to cover the near horizon geometry of rotating black holes and colliding waves with cross polarization. Extreme Kerr and Kerr-Newman geometries are given as examples.

It is believed that matter collisions at speeds almost the speed of light can produce mini black holes even in a laboratory. This possibility directs attentions naturally on collisions of waves in general relativity. Local equivalence (isometry) between two different types of spacetimes, such as black holes (BHs) and colliding plane waves (CPWs) was pointed out first by Chandrasekhar and Xanthopoulos CX [1]. This isometry covered only a portion of the BH geometry, namely region between horizons and the interaction region of CPWs. This concerns only those CPWs which create horizon instead of an all encompassing space-like singularity. Such examples of CPWs are very rare so far. The prototype is the solution of CX which describes collision of impulse gravitational waves accompanied by shock gravitational waves. As a second example we quote colliding electromagnetic (em) shock waves of Bell and Szekeres (BS) [2], which is transformable to the Bertotti-Robinson (BR) solution [3]. This latter is important from its correspondence to conformal field theory and the underlying anti-de Sitter (AdS) structure with a group $SL(2, R) \times SO(3)$. From geometrical point of view BR represents the throat region of an extreme Reissner-Nordstrom BH and its equivalence to colliding em waves is well-known [3].

In this letter we show that the connection between BHs and CPWs [4,5,6] is not restricted only to the local isometry of CX, but seems to be more deep rooted. This we do by showing that the near horizon geometry of the Extreme Kerr and Kerr-Newman BHs are identical with particular CPWs.

The Kerr metric in the Boyer-Linquist coordinates is given by

$$ds^2 = e^{2\nu} d\tilde{t}^2 - e^{2\psi} (d\tilde{\phi} - \omega d\tilde{t})^2 - \rho^2 \left(\frac{d\tilde{r}}{\tilde{\Delta}} + d\theta^2 \right) \quad (1)$$

where

$$\begin{aligned} e^{2\nu} &= \frac{\tilde{\Delta} \rho^2}{(\tilde{r}^2 + a^2)^2 - \tilde{\Delta} a^2 \sin^2 \theta} \\ e^{2(\psi+\nu)} &= \tilde{\Delta} \sin^2 \theta \\ \omega e^{-2\nu} &= \frac{2Ma\tilde{r}}{\tilde{\Delta} \rho^2} \\ \rho^2 &= \tilde{r}^2 + a^2 \cos^2 \theta \\ \tilde{\Delta} &= (\tilde{r} - r_+)(\tilde{r} - r_-) \\ r_+ &= M + \sqrt{M^2 - a^2} \\ r_- &= M - \sqrt{M^2 - a^2} \end{aligned} \quad (2)$$

In the extreme case we have $M = a$, so that angular momentum is $J = M^2$, the horizon area becomes $A = 8\pi M^2$ while $\tilde{\Delta} = (\tilde{r} - M)^2$. The near horizon geometry is described by setting

$$\begin{aligned}
\tilde{r} &= M + \lambda r \\
\tilde{t} &= \frac{t}{\lambda} \\
\tilde{\phi} &= \phi + \frac{t}{2M\lambda}
\end{aligned} \tag{3}$$

and taking the limit $\lambda \longrightarrow 0$. One obtains (after shifting also $r \longrightarrow \frac{1}{r}$) the geometry [7]

$$ds^2 = \frac{(1 + \cos^2 \theta)}{2} \left[\left(\frac{dt}{r_0 r} \right)^2 - \left(\frac{r_0 dr}{r} \right)^2 - (r_0 d\theta)^2 \right] - \frac{2(r_0 \sin \theta)^2}{(1 + \cos^2 \theta)} \left(d\phi - \frac{dt}{rr_0^2} \right)^2 \tag{4}$$

in which $r_0^2 = 2M^2$. We set further, for simplicity $r_0^2 = 2M^2 = 1$. Although BR metric represents the throat region of an extreme Reissner-Nordstrom BH the metric (4) due to its cross term signals more of a BH property. It can easily be checked that the Killing vector ∂_t becomes spacelike for $\cos 2\theta < 4\sqrt{3} - 7$. We show now that this near horizon geometry of an extreme Kerr BH coincides with the colliding wave metric of CX [1]. The CX metric describing collision of impulsive gravitational waves accompanied by shock waves is given by

$$ds^2 = X \left(\frac{d\eta^2}{\Delta} - \frac{d\mu^2}{\delta} \right) - \Delta \delta \frac{X}{Y} dy^2 - \frac{Y}{X} (dx - q_2 dy)^2 \tag{5}$$

where

$$\begin{aligned}
X &= (1 - p\eta)^2 + q^2\mu^2 \\
Y &= p^2\Delta + q^2\delta \\
q_2 &= \frac{2q(1 - \eta)(p\mu^2 + p\eta + \mu^2 - 1)}{(1 + p)(1 - p^2\eta^2 - q^2\mu^2)} \\
\Delta &= 1 - \eta^2 \\
\delta &= 1 - \mu^2
\end{aligned} \tag{6}$$

in which the constants of second polarization (p, q) are constrained by

$$p^2 + q^2 = 1 \tag{7}$$

We note that the metric function q_2 is determined up to an additive constant. This metric admits a horizon at $\eta = 1$ and is free of curvature singularities. Complete analytic extension of this space time was given in detail by CX [1]. In particular the extended spacetime has timelike singularities on hyperbolic arcs reminiscent of the ring singularity in the Kerr space. Now let us choose $p = 0, q = 1$ so that the metric (5) becomes

$$ds^2 = X \left(\frac{d\eta^2}{\Delta} - \frac{d\mu^2}{\delta} - \Delta dy^2 \right) - \frac{4Y}{X} (d\tilde{x} - \eta dy)^2 \tag{8}$$

where

$$\begin{aligned}
X &= 1 + \mu^2 \\
Y &= 1 - \mu^2
\end{aligned} \tag{9}$$

and we have introduced the new coordinate

$$d\tilde{x} = \frac{1}{2}dx + dy$$

Next we apply the transformation

$$\begin{aligned}\eta &= \frac{1}{2r} (r^2 - t^2 + 1) \\ \tanh y &= \frac{1}{2t} (r^2 - t^2 - 1) \\ \mu &= \cos \theta \\ \tilde{x} &= \phi - \frac{1}{2} \ln \left[\frac{(r+t)^2 - 1}{(r-t)^2 - 1} \right]\end{aligned}\tag{10}$$

which results in

$$ds^2 = (1 + \cos^2 \theta) \left(\frac{dt^2 - dr^2}{r^2} - d\theta^2 \right) - \frac{4 \sin^2 \theta}{1 + \cos^2 \theta} \left(d\phi - \frac{1}{r} dt \right)^2\tag{11}$$

This metric is (up to a scale factor of $\frac{1}{2}$) the line element (4) for the geometry of near horizon limit of an extreme Kerr metric. We note that by scaling an overall constant Q^2 can be introduced in front of this metric. This amounts, in the transformation (10) to taking $\tilde{x} \rightarrow Q\tilde{x}$ and $y \rightarrow Qy$. We add also that the choice $p = 0, q = 1$ pushes the timelike singularities on hyperbolic arcs to infinity [1].

The Kerr-Newman metric has the same form as (1) except that $\tilde{\Delta} = \tilde{r}^2 - 2M\tilde{r} + a^2 + q^2$, where q is the electric charge and the $2M\tilde{r}$ factor in ω is replaced by $\tilde{r}^2 + a^2 - \tilde{\Delta}$. The extreme Kerr-Newman case corresponds to $M^2 = a^2 + q^2$ and the horizon is at $\tilde{r} = M$ with area $4\pi(M^2 + a^2)$. Expanding ω to first order in $\tilde{r} - M$ one obtains

$$\begin{aligned}\omega &= \frac{a}{r_0^2} - \frac{2Ma}{r_0^4} (\tilde{r} - M) \\ r_0^2 &= M^2 + a^2\end{aligned}\tag{12}$$

the near horizon limit takes the form [7] (with $r \rightarrow \frac{1}{r}$)

$$\begin{aligned}ds^2 &= \left(1 - \frac{a^2 \sin^2 \theta}{r_0^2} \right) \left[\left(\frac{dt}{r_0 r} \right)^2 - \left(\frac{r_0 dr}{r} \right)^2 - (r_0 d\theta)^2 \right] \\ &\quad - r_0^2 \sin^2 \theta \left(1 - \frac{a^2 \sin^2 \theta}{r_0^2} \right)^{-1} \left(d\phi + \frac{2Ma}{rr_0^4} dt \right)^2\end{aligned}\tag{13}$$

We proceed now to show that this corresponds to a metric of colliding cross polarized em shock waves [8,9]. In the notation of CX this metric is given by

$$ds^2 = F \left(\frac{d\eta^2}{\Delta} - \frac{d\mu^2}{\delta} \right) - \Delta \delta dy^2 - \frac{\delta}{F} (dx - q_0 \eta dy)^2\tag{14}$$

where

$$\begin{aligned}
2F &= 1 + p_0 + (p_0 - 1) \mu^2 \\
p_0 &= \sqrt{1 + q_0^2} = \text{const.}
\end{aligned}
\tag{15}$$

In terms of the null coordinates (u, v) we have $\eta = \sin(au + bv)$ and $\mu = \sin(au - bv)$, where (a, b) are the constant em parameters. For $q_0 = 0$ ($p_0 = 1$) this solution reduces to the BS solution of colliding em waves with linear polarization. Applying a similar transformation as (10) (with $\tilde{x} = x$ now) casts our metric into

$$ds^2 = F(\theta) \left(\frac{dt^2 - dr^2}{r^2} - d\theta^2 \right) - \frac{\sin^2 \theta}{F(\theta)} \left(d\phi - \frac{dt}{r} \right)^2
\tag{16}$$

where

$$2F(\theta) = 1 + p_0 + (p_0 - 1) \cos^2 \theta
\tag{17}$$

Comparing the metrics (13) and (16) we observe that upon choosing $q_0 = -Ma$ with $M^2 = a^2 + 2$ (which corresponds to a charge of $\sqrt{2}$) and scalings $\phi \longrightarrow r_0^2 \phi$, $t \longrightarrow \frac{t}{r_0^2}$ they become identical. This proves that the near horizon geometry of an extreme Kerr-Newman BH is obtained as a result of two colliding em waves with cross polarization.

A general proof, namely, that the extreme geometry of any BH is represented by a CPW remains open. Generic representations of both classes (of BHs and CPWs) may provide a general proof. We are satisfied here only with the two simplest but most important cases. A general proof on the other hand is reminiscent of a theorem by Penrose [10]: Any spacetime admits a plane wave limit. Here it is not any spacetime but it is only the near horizon geometry of an extreme BH. And it is not a plane wave limit but it is a colliding wave limit

Acknowledgement 1 *I wish to thank Ozay Gurtug for helpful discussions.*

REFERENCES

- [1] S. Chandrasekhar and B.C. Xanthopoulos, Proc. R. Soc. London A408, 175 (1986).
- [2] P. Bell and P. Szekeres, Gen. Rel. Grav. 5, 275 (1974).
- [3] J.B.Griffiths, Colliding Plane Waves in General Relativity (Oxford University Press, Oxford,1991) p.138
- [4] S.A. Hayward, Class. Quantum Grav. 6, 1021 (1989).
- [5] U. Yurtsever, Phys. Rev. D37, 2790 (1988).
- [6] P. Schwarz, Phys. Rev. D56, 7833 (1997).
- [7] J. Bardeen and G.T. Horowitz, Phys. Rev. D60, 104030 (1999).
- [8] M. Halilsoy, Phys. Rev. D37, 2121 (1988).
- [9] M. Halilsoy, J. Math. Phys. 31, 2694 (1990).
- [10] R. Penrose, "Any spacetime has a plane wave as a limit", in Differential Geometry, pp271 Reidel, Dordrecht, 1976.